


Coronavirus FAQ for Shelter & Drop-In Settings

A resource for people working in high-risk community settings.







COVID-19 is spread by droplets; it can come from a cough or sneeze, someone's hand, or even un unclean surface.



Disinfect common areas more frequently. Use alcohol solutions with >70% alcohol, or diluted household bleach.



The 3 hallmark symptoms of COVID-19 are:

- **1. Fever (≥38°C)**
- 2. Cough (Usually Dry)
- 3. Shortness of Breath



Wear a facemask to cover your coughs and sneezes.

Ask symptomatic clients to wear a mask, and do frequent hand washing.



Social distancing means **6 feet or 2 metes.**

Have clients sleeping head-to-toe.



Let your manager know if you are at higher risk for health complications related to COVID-19.



Soap and water is the **most effective** hand cleanser.

If unavailable, hand sanitizer with **>60% Alcohol**



Designate someone to **monitor** clients for symptoms.

Reassure clients that you are taking **precautions** to keep them safe.



Frequently Asked Questions



A resource for people working in high-risk community settings.

n	ue	cti		nc
U	uc	3 L I	U	113
~~				

Answers

What is the personal protective equipment overdose response during the COVID-19 outbreak?



Your naloxone kits contain **one-way pocket valve** masks which will provide protection.

The use of BVM increases the risk of transmission; it does not prevent it. Therefore currently **we do not recommend** using BVM during an OD response.

What is the proper technique for disinfecting surfaces?



Clean the surface with soap and water first, then apply approved disinfectant. Let sit for 10 minutes before wiping.

A simple 'spray & wipe' won't work!

What should I do if we don't have enough supplies?



Contact you local **BC Housing** representative for help with:

-Soap - Alcohol based hand sanitizer - Tissues -Disinfecting wipes - Trash baskets - Gloves -Disposable masks - Paper plates/cutlery



Should I wear a mask all the time? Should I wear two masks?



Masks are in short supply and should be used during periods of higher risk wearing two masks can actually **increase risk.**